

February 1, 2008

Algebraic treatment of the confluent Natanzon potentials

M. T. Chefrour

Département de Physique, Faculté des Sciences,
Université Badji Mokhtar, Annaba, Algérie.

L. Chetouani and L. Guechi

Département de Physique, Faculté des Sciences,
Université Mentouri, Route d'Ain El Bey,
Constantine, Algérie.

Abstract

Using the $so(2, 1)$ Lie algebra and the Baker, Campbell and Hausdorff formulas, the Green's function for the class of the confluent Natanzon potentials is constructed straightforwardly. The bound state energy spectrum is then determined. Eventually, the three-dimensional harmonic potential, the three-dimensional Coulomb potential and the Morse potential may all be considered as particular cases.

PACS 03.65-Quantum theory ; quantum mechanics.

PACS 03.65.Fd -Algebraic methods.

PACS 02.20.+b -Group theory.

typescript using Latex (version 2.0).

In this paper, we shall want to present an algebraic treatment of the bound state problem for the class of the confluent Natanzon potentials defined by

$$V(r) = \frac{g_2 h^2 + g_1 h + \eta}{R} + \frac{\sigma_1 h - \sigma_2 h^2}{R^2} - \frac{5}{4} \frac{\Delta h^2}{R^3}, \quad (1)$$

where $\Delta = \sigma_1^2 - 4\sigma_2 c_0$ and $g_1, g_2, \sigma_1, \sigma_2, c_0$ and η are dimensionless parameters. The function $h = h(r)$ is defined implicitly by the differential equation

$$\frac{dh}{dr} = \dot{h} = \frac{2h}{\sqrt{R(r)}}, \quad (2)$$

where

$$R(r) = \sigma_2 h^2 + \sigma_1 h + c_0. \quad (3)$$

These potentials, originally introduced by Natanzon, are interesting in the sense that the Schrödinger equation can be reduced to the confluent hypergeometric form and they include, as special cases, the radial harmonic oscillator for $\sigma_2 = c_0 = 0$, the radial Coulomb potential for $\sigma_1 = c_0 = 0$ and the Morse potential for $\sigma_2 = \sigma_1 = 0$. These confluent Natanzon potentials have been recently the object of several studies. More specifically, Natanzon¹ has obtained, in a Schrödinger equation approach, the discrete energy spectrum, the unnormalized wave functions and has derived the Green's function. Cooper and al² have used the supersymmetric quantum mechanics as an algebraic method to discuss these potentials. Furthermore, the spectrum of these potentials has been re-derived by Cordero and Salamó³, still in a Schrödinger approach, by making use a particular realization of the $so(2, 1)$ algebraic method which is called spectrum generating algebra (SGA)⁴. Finally, this class of the confluent Natanzon potentials has been solved very recently by Grosche⁵ in the framework of a Feynman path integral approach.

The Milshtein and Strakhovenko variant (MS) of the $so(2, 1)$ algebraic approach⁶ provides an elegant and straightforward way of solving the problem for $V(r)$. As we shall see, this approach consists of constructing the Green's function by expressing the resolvent operator for this system in the form of a linear combination of an appropriate realization of $SO(2, 1)$ generators. This form is then written in the Schwinger's integral representation and the action of the resolvent operator is obtained by means of two Baker, Campbell and Hausdorff formulas.

Let $G(r, r'; E)$ be the Green's function associated to the potentials $U(r) = \frac{\hbar^2}{2m} V(r)$, which is solution of the differential equation

$$(H - E) G(r, r'; E) = \frac{\hbar}{i} \delta(r - r'), \quad (4)$$

where

$$H = -\frac{\hbar^2}{2m} \frac{d^2}{dr^2} + U(r) \quad (5)$$

is the Hamiltonian of the system and E its energy.

In terms of the function $h(r)$ and by taking into account the relation (2), we can write the resolvent operator $H - E$ as follows:

$$\begin{aligned} H - E = & \frac{\hbar^2}{2m} \left\{ -\frac{d^2}{dr^2} + \frac{1}{4} \left(3\frac{\ddot{h}^2}{\dot{h}^2} - 2\frac{\ddot{h}}{\dot{h}} \right) \right. \\ & \left. + \frac{\dot{h}^2}{4h^2} [(g_2 - \sigma_2\epsilon)h^2 + (g_1 - \sigma_1\epsilon)h + (\eta - 1 - c_0\epsilon)] \right\}, \quad (6) \end{aligned}$$

with $E = \frac{\hbar^2}{2m}\epsilon$.

The point canonical transformation $\xi = h(r)$ leads to

$$(\widetilde{H} - \widetilde{E}) \widetilde{G}(\xi, \xi'; \widetilde{E}) = \frac{\hbar}{i} \xi \delta(\xi - \xi'), \quad (7)$$

where

$$\widetilde{G}(\xi, \xi'; \widetilde{E}) = \sqrt{\dot{h}(r) \dot{h}(r')} G(r, r'; E). \quad (8)$$

Then, the dynamics of the physical system is governed by the new Hamiltonian

$$\widetilde{H} = \frac{\hbar^2}{2m} \left(-\xi \frac{d^2}{d\xi^2} + \frac{\eta - 1 - c_0\epsilon}{4\xi} + \frac{g_2 - \sigma_2\epsilon}{4} \xi \right), \quad (9)$$

and

$$\widetilde{E} = -\frac{\hbar^2}{8m} (g_1 - \sigma_1\epsilon) \quad (10)$$

is the pseudo-energy.

To see that one can introduce the three following generators

$$T_1(x) = -\frac{\hbar^2}{2M} \left(\frac{d^2}{dx^2} - \frac{\mu(\mu-1)}{x^2} \right), T_2(x) = -\frac{i}{2} \left(x \frac{d}{dx} + \frac{1}{2} \right), T_3(x) = \frac{M}{4\hbar^2} x^2, \quad (11)$$

satisfying the $so(2, 1)$ Lie algebra⁷

$$[T_1, T_2] = -iT_1, \quad [T_2, T_3] = -iT_3, \quad [T_1, T_3] = -iT_2, \quad (12)$$

it is convenient to perform the change of variable $\xi = x^2$ and redefine the Green's function as

$$\tilde{G}(\xi, \xi'; \tilde{E}) = \sqrt{xx'} G(x, x'; \tilde{E}). \quad (13)$$

Under this change the differential equation (7) becomes

$$\left(T_1(x) + 2\hbar^2 \tilde{\omega}^2 T_3(x) - \tilde{E} \right) G(x, x'; \tilde{E}) = \frac{\hbar}{i} \delta(x - x'), \quad (14)$$

where

$$\tilde{\omega} = \frac{\hbar}{M} (g_2 - \sigma_2 \epsilon)^{\frac{1}{2}} \quad (15)$$

with $M = 4m$.

Being a linear combination of the generators T_i , the operator $\tilde{H}$ shows, as expected, a dynamical symmetry $so(2, 1)$. By using the Schwinger's integral representation⁸, the solution of the differential equation (14) can be written as follows:

$$G(x, x'; \tilde{E}) = \int_0^\infty dS \exp \left[\frac{i}{\hbar} (\tilde{E} + i0) S \right] K(x, x'; S), \quad (16)$$

where

$$K(x, x'; S) = \exp \left\{ -\frac{iS}{\hbar} [T_1(x) + 2\hbar^2 \tilde{\omega}^2 T_3(x)] \right\} \delta(x - x'). \quad (17)$$

The calculation of this kernel is based upon the use of two Baker, Campbell- and Hausdorff formulas⁷

$$\exp \left\{ -\frac{iS}{\hbar} [T_1 + 2\hbar^2 \tilde{\omega}^2 T_3] \right\} = \exp(-iaT_3) \exp(-ibT_2) \exp(-icT_1), \quad (18)$$

where

$$a = 2\hbar\tilde{\omega} \tan(\tilde{\omega}S), \quad b = 2 \ln [\cos(\tilde{\omega}S)], \quad c = \frac{1}{\hbar\tilde{\omega}} \tan(\tilde{\omega}S), \quad (19)$$

and

$$\exp(-i\alpha T_3) \exp(-i\beta T_2) \exp(-i\gamma T_1) = \exp(-icT_1) \exp(\tau T_3), \quad (20)$$

with

$$\alpha = \frac{i\tau}{1 - i\tau c/2}, \quad \beta = 2 \ln \left(1 - \frac{i\tau c}{2} \right), \quad \gamma = \frac{c}{1 - i\tau c/2}. \quad (21)$$

It is to be noted that formulas (18) and (20) can be easily checked using the finite representation of $so(2, 1)$ Lie algebra where the T_i operators are defined in terms of σ_i Pauli matrices

$$T_1 = \frac{\sigma_1 - i\sigma_2}{2\sqrt{2}}, \quad T_2 = -\frac{i\sigma_3}{2}, \quad T_3 = \frac{\sigma_1 + i\sigma_2}{2\sqrt{2}}. \quad (22)$$

Here, we also have to use the Laplace transform of the Dirac distribution

$$\delta(x - x') = \frac{M}{2\hbar^2} \frac{x^\mu x'^{1-\mu}}{2i\pi} \int_{-i\infty+\delta}^{i\infty+\delta} d\tau \exp \left[\frac{M}{4\hbar^2} (x^2 - x'^2) \tau \right]; \delta \prec 0, \quad (23)$$

in order to obtain a manageable result as follows:

$$\exp(-i\gamma T_1) x^\mu = \left[1 - i\gamma T_1 + \frac{1}{2!} (-i\gamma T_1)^2 + \dots \right] x^\mu = x^\mu. \quad (24)$$

Using relations (23), (18) and (20), the kernel (17) can now be written

$$\begin{aligned} K(x, x'; S) &= \frac{M}{2\hbar^2} x'^{1-\mu} \exp(-iaT_3) \exp(-ibT_2) \frac{1}{2i\pi} \int_{-i\infty+\delta}^{i\infty+\delta} d\tau \\ &\quad \times \exp \left(-\frac{M}{4\hbar^2} x'^2 \tau \right) \exp(-icT_1) \exp \left(\frac{M}{4\hbar^2} x^2 \tau \right) x^\mu \\ &= \frac{M}{2\hbar^2} x'^{1-\mu} \exp(-iaT_3) \exp(-ibT_2) \frac{1}{2i\pi} \int_{-i\infty+\delta}^{i\infty+\delta} d\tau \\ &\quad \times \exp \left(-\frac{M}{4\hbar^2} x'^2 \tau \right) \exp(-icT_1) \exp(\tau T_3) x^\mu \\ &= \frac{M}{2\hbar^2} x'^{1-\mu} \exp(-iaT_3) \exp(-ibT_2) \frac{1}{2i\pi} \int_{-i\infty+\delta}^{i\infty+\delta} d\tau \\ &\quad \times \exp \left(-\frac{M}{4\hbar^2} x'^2 \tau \right) \exp(-i\alpha T_3) \exp(-i\beta T_2) \exp(-i\gamma T_1) x^\mu \\ &= \frac{M}{2\hbar^2} x'^{1-\mu} \exp(-iaT_3) \exp(-ibT_2) x^\mu \frac{1}{2i\pi} \\ &\quad \times \int_{-i\infty+\delta}^{i\infty+\delta} d\tau \frac{\exp \left\{ \frac{M}{2\hbar^2} \tau \left(-\frac{x'^2}{2} + \frac{x^2}{2-i\tau c} \right) \right\}}{\left(1 - \frac{i\tau c}{2} \right)^{\mu+\frac{1}{2}}} \\ &= \frac{M}{2\hbar^2} x'^{1-\mu} \exp(-iaT_3) \exp(-ibT_2) x^\mu \exp \left(\frac{iMx^2}{2\hbar^2 c} \right) \\ &\quad \times \int_{-i\infty+\delta}^{i\infty+\delta} d\tau \frac{\exp \left(-\frac{Mx'^2}{4\hbar^2} \tau \right) \exp \left(\frac{Mx^2}{\hbar^2 c^2} \frac{1}{\tau+2i/c} \right)}{\left(1 - \frac{i\tau c}{2} \right)^{\mu+\frac{1}{2}}}, \end{aligned} \quad (25a)$$

where formula^{9,10}

$$\exp(-i\beta T_2) f(x) = \exp \left(-\frac{\beta}{4} \right) f \left(e^{-\frac{\beta}{2}} x \right) \quad (26)$$

has also been used.

The integral can be calculated thanks to the residue theorem after the $\exp\left(\frac{Mx^2}{\hbar^2 c^2} \frac{1}{\tau+2i/c}\right)$ series has been effected. Hence we obtain

$$\begin{aligned}
K(x, x'; S) &= \frac{M}{i\hbar^2 c} \exp(-iaT_3) \exp(-ibT_2) \sqrt{xx'} \exp\left[\frac{iM}{2\hbar^2 c}(x^2 + x'^2)\right] \\
&\quad \times I_{\mu-\frac{1}{2}}\left(\frac{Mxx'}{i\hbar^2 c}\right) \\
&= \frac{M}{i\hbar^2 c} \exp\left(-\frac{iaM}{4\hbar^2} x^2\right) \exp\left(-\frac{b}{2}\right) \sqrt{xx'} \\
&\quad \times \exp\left[\frac{iM}{2\hbar^2 c}(e^{-b}x^2 + x'^2)\right] I_{\mu-\frac{1}{2}}\left(\frac{Mxx'}{i\hbar^2 c e^{\frac{b}{2}}}\right) \\
&= \frac{M\tilde{\omega}}{i\hbar \sin(\tilde{\omega}S)} \sqrt{xx'} \exp\left[\frac{iM\tilde{\omega}}{2\hbar}(x^2 + x'^2) \cot(\tilde{\omega}S)\right] \\
&\quad \times I_{\mu-\frac{1}{2}}\left(\frac{M\tilde{\omega}xx'}{i\hbar \sin(\tilde{\omega}S)}\right), \tag{27}
\end{aligned}$$

where

$$\mu = \frac{1}{2} + \sqrt{\eta - c_0\epsilon}, \tag{28}$$

and $I_{\mu-\frac{1}{2}}(x)$ is the modified Bessel function.

After inserting (27) into (16) and performing the integration over the time variable S with the help of the following formula (see Gradshtein and Ryzhik¹¹, p.729, eq. (6.669.4))

$$\begin{aligned}
&\int_0^\infty dq \frac{e^{-2pq}}{\sinh q} \exp\left[-\frac{1}{2}(x+y) \coth q\right] I_{2\gamma}\left(\frac{\sqrt{xy}}{\sinh q}\right) \\
&= \frac{\Gamma\left(p+\gamma+\frac{1}{2}\right)}{\Gamma(2\gamma+1)\sqrt{xy}} M_{-p,\gamma}(x) W_{-p,\gamma}(y), \tag{29}
\end{aligned}$$

valid for $Re\left(p+\gamma+\frac{1}{2}\right) > 0$, $Re(\gamma) > 0$ and $y > x$, where $M_{-p,\gamma}(x)$ and $W_{-p,\gamma}(y)$ are the Whittaker functions, the Green's function (16) becomes

$$G(x, x'; \tilde{E}) = \frac{\Gamma\left(p+\frac{\mu}{2}+\frac{1}{4}\right)}{2i\tilde{\omega}\Gamma(\mu+\frac{1}{2})\sqrt{xx'}} M_{-p,\frac{\mu}{2}-\frac{1}{4}}\left(\frac{M\tilde{\omega}}{\hbar}x'^2\right) W_{-p,\frac{\mu}{2}-\frac{1}{4}}\left(\frac{M\tilde{\omega}}{\hbar}x^2\right), \tag{30}$$

where $p = -\frac{\tilde{E}}{2\hbar\omega}$ and $x > x'$.

Eventually, by taking into account eqs. (30), (13), (8), (2) and the successive transformations $\xi = h(r) = x^2$, the Green's function associated with the confluent Natanzon potentials may be written:

$$G(r, r'; E) = \frac{\Gamma\left(-\frac{\tilde{E}}{2\hbar\tilde{\omega}} + \frac{\mu}{2} + \frac{1}{4}\right)}{4i\tilde{\omega}\Gamma(\mu + \frac{1}{2})} \left[\frac{\sqrt{R(r)R(r')}}{h(r)h(r')} \right]^{\frac{1}{2}} M_{-p, \frac{\mu}{2} - \frac{1}{4}}\left(\frac{M\tilde{\omega}}{\hbar}h(r')\right) W_{-p, \frac{\mu}{2} - \frac{1}{4}}\left(\frac{M\tilde{\omega}}{\hbar}h(r)\right), \quad (31)$$

and the bound state energy eigenvalues $E_n = \frac{\hbar^2}{2m}\epsilon_n$ can be obtained from the poles of $\Gamma\left(-\frac{\tilde{E}}{2\hbar\tilde{\omega}} + \frac{\mu}{2} + \frac{1}{4}\right)$ in (31),

$$-\frac{\tilde{E}}{2\hbar\tilde{\omega}} + \frac{\mu}{2} + \frac{1}{4} = -n, \quad n \in \mathbf{N}_0. \quad (32)$$

Upon substitution of (10), (15) and (28) into (32), we see then that

$$\frac{g_1 - \sigma_1\epsilon}{2\sqrt{g_2 - \sigma_2\epsilon}} + \sqrt{\eta - c_0\epsilon} = -(2n + 1) \quad (33)$$

is an equation of fourth degree in ϵ_n which can be solved after some simple manipulations¹².

Many particular cases of the potentials (1) arise when some of its parameters vanish. In the following table, we give the interesting ones, namely, the radial harmonic oscillator, the radial Kepler-Coulomb potential and the Morse potential

σ_1	σ_2	c_0	$h(r)$	$V(r)$
	0	0	r^2/σ_1	$\frac{g_1}{\sigma_1} + \frac{g_2}{\sigma_1^2}r^2 + \frac{\eta-1}{r^2}$
0		0	$2r/\sqrt{\sigma_2}$	$\frac{g_2}{\sigma_2} + \frac{g_1}{2\sqrt{\sigma_2}r} + \frac{\eta-1}{4r^2}$
0	0		$e^{-r/\sqrt{c_0}}$	$\frac{g_2}{c_0}e^{-2r/c_0} + \frac{g_1}{c_0}e^{-r/\sqrt{c_0}} + \frac{\eta}{c_0}$

In conclusion, we have shown that the class of confluent Natanzon potentials can be described by a particular realization of the $so(2, 1)$ which enables us to work out the Green's function. The realization of the $so(2, 1)$ algebra used here differs substantially from that of Cordero and Salamó³. The discrete spectrum is identical with the results obtained by using either the (SGA) method³ or the Feynman path integral approach⁵.

References

- ¹G. N. Natanzon, Teor. Mat. Fiz. **38** (1979) 146.
- ²F. Cooper, J. Ginocchio and A. Kahare, Phys. Rev. **D 36** (1987) 2458.
- ³P. Cordero and S. Salamó, J. Phys. A: Math. Gen. **24** (1991) 5299; Condensed Matter Theories, **7** (1992) 49; Found. Phys. **23** (1993) 675.
- ⁴P. Cordero and G. C. Ghirardi, Nuovo Cimento **A 2** (1971) 217; Fortschr. Phys. **20** (1972) 105.
- ⁵C. Grosche, J. Phys. A: Math. Gen. **29** (1996) L183.
- ⁶A. I. Milshtein and V. M. Strakhovenko, Phys. Lett. **A 90** (1982) 447.
- ⁷B. G. Wybourne, *Classical Groups for Physicists* (Wiley, New York, 1974).
- ⁸J. Schwinger, Phys. Rev. **82** (1951) 664.
- ⁹L. Chetouani, L. Guechi and T. F. Hammann, Helv. Phys. Acta **65** (1992) 1069.
- ¹⁰L. Chetouani, L. Guechi and T. F. Hammann, J. Math. Phys. **33** (1992) 3410.
- ¹¹I. S. Gradshtein and I. M. Ryzhik, *Table of Integrals, Series and Products* (Academic, New York, 1965).
- ¹²I. N. Bronshtein and K. A. Semendyayev, *A Guide Book to Mathematics* (Harri Deutsch, Frankfurt/Main 1973).